

[Web](#) [Images](#) [Videos](#) [Maps](#) [News](#) [Shopping](#) [Gmail](#) [more ▼](#)
[abdullah.kawsar@gmail.com](#) | [Scholar Preferences](#) | [My Account](#) | [Sign out](#)

[Advanced Scholar Search](#)
Scholar
☐ Articles excluding patents

 -
☐ include citations

☐ Create email alert

Results 1 - 20 of about 458. (0.65 sec)

The implementation of synchronizing operations in various environments

H Wettstein - Software: Practice and Experience, 1977 - interscience.wiley.com

... If the **interrupt** prevention is located within the scope of the locking variable then the process ... is apparently the fact that two thresholds are erected in front of the **critical region** which leads to ... If it is not available the **threshold** can be simulated by an arrangement shown in Figure 3 ...

[Cited by 10](#) - [Related articles](#) - [All 2 versions](#)

[PDF] Crossover from superconductivity to a negative magnetoresistance in a Cd-Sb alloy near the localization threshold as the temperature is lowered

[\[PDF\] from issp.ac.ru](#)

VF Gantmakher, VN Zverev, VM Teplinskiĭ... - ... C/C OF PIS'MA V ..., 1992 - issp.ac.ru

... It turns out that in the **critical region** the onset temperature of the S transition, T_n , is ... a correlation between the superconducting response and the conductivity type on the / side of the **threshold**. ... By carrying out the heating in increments, one can **interrupt** this process and obtain, in ...

[Cited by 2](#) - [Related articles](#) - [BL Direct](#) - [All 5 versions](#)

Analysis of phylogenetically reconstructed mutational spectra in human mitochondrial DNA control region

BA Malyarchuk, IB Rogozin, VB Berikov... - Human genetics, 2002 - Springer

... procedure was repeated 10,000 times, a critical value P^* that separates the right **critical region** of the ... sites with 3 or more mutations which was used as the hotspot **threshold** ($Sh=3$... Interestingly, the additional mutations (at positions 16186 and 16192) that **interrupt** the poly-C tract ...

[Cited by 46](#) - [Related articles](#) - [BL Direct](#) - [All 4 versions](#)

Genes and translocations involved in POF

D Schiessinger, L Herrera, L Crispont... - American journal of ..., 2002 - Wiley Online Library

... follicles after menarche and is progressively depleted until it falls below a **threshold** value and ... If ovarian genes are clustered in the **critical region**, every breakpoint in the region should affect ... The notion that translocations **interrupt** genes to cause POF is further weakened by the ...

[Cited by 83](#) - [Related articles](#) - [BL Direct](#) - [All 4 versions](#)

Subacute electrical stimulation of the hippocampus blocks intractable temporal lobe seizures and paroxysmal EEG activities

M Velasco, F Velasco, AI Velasco, B Boileaga... - Epilepsia, 2000 - Wiley Online Library

... stimulation with rectangular pulses **interrupted** only 1 h/day while EEG recordings were taken. Individual pulses consisted of biphasic, Lilly wave pulses 130 per second in frequency, 450 ps in duration, 200-400 pA in amplitude. Amplitude was adjusted for **threshold** or slightly ...

[Cited by 132](#) - [Related articles](#) - [BL Direct](#) - [All 4 versions](#)

A scheduling scheme for network saturated NT multiprocessors

[\[PDF\] from psu.edu](#)

JS Hansen... - Proceedings of the USENIX Windows NT Workshop, 1997 - unix.org

... it possible to specify that a certain amount of the workload must be performed in a **critical region**. ... sharing data protected by locks, the effects of the user thread starvation caused by **interrupt**-routines may ... The fixed **threshold** used to shift to polling is a problem in the case where ...

[Cited by 10](#) - [Related articles](#) - [All 12 versions](#)

Development of the concurrent process oriented language'COL'

M Mitsuishi, I Shimoyama... - Advanced Robotics, 1987 - ingentaconnect.com

... control, (c) priority control, (d) shared variable management, (e) real-time operation, (f) I/O handling, (g) **interrupt** handling, and (h) process state control. ... The **critical region** for mutual exclusion is described using 'region' as follows: ...

[Related articles](#) - [All 5 versions](#)

Color-induced transition to a nonconventional diffusion regime

GP Tsironis... - Physical review letters, 1988 - APS

... by construction, the initial population at $x=0$, $c(x;0)$, vanishes throughout the whole **critical region** $T_z r$... Figure 1 shows that $r(O,t)$ is always semidefinite positive even beyond the **threshold** T_r , where ... T/T_0) on I/I_0 , does not provide the correct slope ' and does not **interrupt** this as ...

[Cited by 12](#) - [Related articles](#) - [All 5 versions](#)

Electrophysiological and behavioral effects of blockade of the nonspecific thalamo-cortical system

JE Skinner... - Brain Research, 1967 - Elsevier

... of varying the temperature of the cryo-probe tip when in the **critical region** of ITP. ... are enhanced when any part of the total nonspecific thalamo-cortical system is **interrupted** or blocked ... of the cortex, corresponding to an inhibitory state in that region and a high **threshold** of excitability ...

[Cited by 84](#) - [Related articles](#) - [All 2 versions](#)

Role of midbrain central gray in pain-induced defensive boxing of rats

MA Edwards... - Physiology & Behavior, 1974 - Elsevier

... lesions which extended in anterior, posterior, or ventral directions from the **critical region** (Fig. ... became slightly less reactive on their second test (mean change in **threshold**: +13.0 ... Similarly,

these lesions **interrupted** the major descending pathways from the forebrain to the central ...

[Cited by 39](#) - [Related articles](#) - [All 4 versions](#)

Comparison of phenotype between patients with Prader-Willi syndrome due to deletion 15q and uniparental disomy 15

SB Cassidy, M Forsythe, S Heeger... - Journal of Medical ..., 1997 - interscience.wiley.com
... using highly polymorphic microsatellite markers both within the Prader-Willi Syndrome **Critical Region** (15q11-q13 ... is that one or both of the breakpoints for the deletion **interrupt** a gene ... insufficient for clear recommendations, but it would appear that the **threshold** for clinical ...

[Cited by 118](#) - [Related articles](#) - [BL Direct](#) - [All 4 versions](#)

Percolation in short fibres epoxy resin composites: Conductivity behavior and finite size effects near threshold

F Carmona, P Prud'homme... - Solid State Communications, 1984 - Elsevier
... conductivity exponent the reason being that the experiments have been performed outside the **critical region**. ... In the vicinity of the percolation **threshold** where $\sim \sim 1@- @*1-v$ this leads ... Full, dashed and **interrupted** lines give hypothetical power law dependences with $u = 1, 0.88$...

[Cited by 45](#) - [Related articles](#) - [All 4 versions](#)

Catheter ablation of the mitral isthmus for ventricular tachycardia associated with inferior infarction

[\[HTML\] from ahajournals.org](#)

DJ Wilber, DE Kopp, DO Glascock, N Dennis... - Circulation, 1995 - Am Heart Assoc
... Conclusions The mitral isthmus contains a **critical region** of slow conduction in some patients with ... using constant current with a 2-ms pulse duration at twice diastolic **threshold**. ... narrow dimensions, may have been simply the most vulnerable point to **interrupt** this circumferential ...

[Cited by 111](#) - [Related articles](#) - [BL Direct](#) - [All 5 versions](#)

Brain stimulation for neurological and psychiatric disorders, current status and future direction

[\[HTML\] from aspetjournals.org](#)

JY Chang - Journal of Pharmacology and Experimental ..., 2004 - ASPET
... Anterior and centromedian thalamic stimulations may **interrupt** epileptiform activity relaying to the cortices thus ... The thalamus is the **critical region** for relaying neural signals to the cerebral cortex and ... et al., (1982) pointed out that naloxone elevated the stimulation **threshold** in the ...

[Cited by 16](#) - [Related articles](#) - [BL Direct](#) - [All 5 versions](#)

Conduction in anisotropic disordered systems: Effective-medium theory

J Bernasconi - Physical Review B, 1974 - APS
... 4 It has been successfully applied to percolation models (outside the **threshold** region) and ... on networks with 20x20 nodes is remarkably good, except in a **critical region** near p ... This boundedness of the anisotropy Q has important consequences for the **interrupted** strand model ...

[Cited by 42](#) - [Related articles](#) - [All 2 versions](#)

Neurons of rostral ventrolateral medulla mediate somatic pressor reflex

Rt Stornetta, SF Morrison... - American Journal of ..., 1989 - Am Physiological Soc
... that it is mediated by spinoreticular afferents traveling in the contralateral spinal cord, and that the CI adrenergic area of the RVL is a **critical region** for the ... stimulation (stimulus trace) is a 5-s train of square-wave pulses (1.0 ms, 20 Hz) at a stimulus current 5 times **threshold** (see ...

[Cited by 31](#) - [Related articles](#) - [All 3 versions](#)

Toward a molecular understanding of congenital heart disease

[\[HTML\] from ahajournals.org](#)

RM Payne, MC Johnson, JW Grant... - Circulation, 1995 - Am Heart Assoc
... resultant mutant phenotype will provide clues to the normal function of the **interrupted** gene ... In the **threshold** model, which some have proposed to explain human CHD, 2 multiple ... combined with the trisomy 16 (Ts16) mouse model have narrowed the **critical region** for endocardial ...

[Cited by 111](#) - [Related articles](#) - [BL Direct](#) - [All 5 versions](#)

Phase transitions in relational learning

[\[PDF\] from uniol.it](#)

A Giordana... - Machine Learning, 2000 - Springer
... From figures 8(a)-(d) we would predict that formula ϕ_1 should be easy to match for all the examples, whereas ϕ_2 is likely to require a high computational cost to be matched, because most examples lay in the **critical region**. ...

[Cited by 56](#) - [Related articles](#) - [BL Direct](#) - [All 5 versions](#)

[HTML] Disruption of a novel member of a sodium/hydrogen exchanger family and DOCK3 is associated with an attention deficit hyperactivity disorder-like phenotype

[\[HTML\] from bmj.com](#)

MG De Silva, K Elliott, HH Dahl... - Journal of medical ..., 2003 - jmg.bmj.com
... assembled using computer analyses of EST markers along the BAC clones in the **critical region** and sequence ... The gene **interrupted** at the p arm, DOCK3, is a member of the CDM family ... Although the lod score did not exceed the QTL **threshold** for significance, its location within ...

[Cited by 39](#) - [Related articles](#) - [BL Direct](#) - [All 3 versions](#)

Approximate analysis of distributed semi-hard real-time systems

D Mañnesco - Automatic Control, IEEE Transactions on, 2002 - ieeexplore.ieee.org
... and S. B. Gershwin, "The uncertainty **threshold** principle: Some fundamental limitations of optimal decision making under ... CS, is a senice center corresponding to execution in **critical region** i. ... A process with a higher priority cannot **interrupt** the execution of a process with lower ...

[Cited by 7](#) - [Related articles](#) - [All 3 versions](#)

☒ [Create email alert](#)